



**State of the Lake
Environment Report
2011**

Bennett Lake



Mississippi Valley Conservation

Why Watershed Watch?

The Ribbon of Life is the shallow waters and first 10 metres of shore land around our lakes where the land meets the water. This ribbon—where land meets the water—is where much of the lake life is born, raised and fed. Many landowners, unaware of the importance of this area, have cleared the shorelines of native vegetation and replaced it with lawns, non-native ornamental vegetation, retaining walls and boathouses. This has had a negative effect on fish and wildlife habitat and water quality. Natural vegetation retained or restored along the shoreline helps prevent erosion and improves water quality by binding nutrients before they can enter the lake.

Mississippi Valley Conservation (MVC) has long recognized the recreational and aesthetic value of lakes within the watershed and is committed to maintaining and protecting water quality and fish habitat. MVC has joined together with volunteer lake stewards throughout the watershed to take steps to protect and restore water quality by launching the Watershed Watch program in 1998. Watershed Watch is an environmental monitoring and awareness program. The objectives of the program are to collect reliable environmental data to document current water quality conditions and use the data as an essential educational tool to encourage residents to adopt sound stewardship practices aimed at preserving and protecting water quality. Together we will encourage and assist shoreline residents, both seasonal and permanent, to become personal stewards of their lake by taking an active role in restoring and enhancing their shoreline to maintain water quality and a healthy lake environment.



Easy steps to improve water quality

Build at least 30 metres (100 feet) away from the shoreline

Keep your lot well treed

Preserve or replant native vegetation along the shoreline

Pump out your septic tank every three to five years

Reduce water usage

Use phosphate free soaps and detergents

Keep the size of your lawn to a minimum

Bennett Lake

This lake was last surveyed using imperial measurements, therefore the original bathymetric (depth) measurements are recorded in feet rather than metres.

3.28 ft = 1 m 1ft = 0.3048 m

This map is intended for illustration only; it should not be used as a navigation guide.



The Secchi Disk

- It's a round disk that is lowered into a lake to visually measure water clarity.
- The depth at which the disk disappears indicates the level of nutrients and algae growth.
- The higher the reading, the clearer the lake. The more nutrients that run into the lake, the more algae grows, causing reduced water clarity.

Depth: Your lake is likely...

- | | |
|----------|----------------------------------|
| 0 - 3 m. | enriched, eutrophic |
| 3 - 5 m. | moderately enriched, mesotrophic |
| 5+ m. | unenriched, oligotrophic |



Most Common Species of Fish:

- Largemouth Bass
- Smallmouth Bass
- Rock Bass
- Walleye
- Northern Pike
- White Sucker
- Northern Redhorse Sucker
- Yellow Perch
- Burbot
- Brown Bullhead
- Pumpkin Seed

Remember:
Use non-lead sinkers to protect the health of the fish and this lake.



Bennett Lake

Located in the Townships of Bathurst Burgees and Sherbrooke.
 Elevation of 158 metres above sea level.
 Lake perimeter is 35 kilometres
 Deepest point is 12.2 metres.
 Warm water fishery including; Northern Pike, Small and Largemouth Bass,
 Yellow Perch and Rock Bass.
 According to Tay Valley Township as of 2006 there were 205 properties with
 lake frontage.

How Does Bennett Lake Measure Up?

1980 – 2006 Water Quality Results – South Basin


Sample Year Mean	Secchi Disc Depth (Metres)	Total Phosphorus Euphotic Zone (Micrograms/litre)	Total Phosphorus 1 Metre off Bottom (Micrograms/litre)	Chlorophyll <u>a</u> Composite (Micrograms/litre)
**1975	3	22	20	6.94
**1980	3.4	13.2	230	5.49
1997	5	18	n/a	n/a
2001	3.5	11.6	138.5	5.73
*2006	4.4	11.3	28.7	2.73
2011	4.2	19.3	57.7	5.2
N Mean	6 3.9	6 15.9	5 95.0	5 5.2
Standard Deviation	0.744087808	4.474818432	88.78235748	1.540444741

2001 - 2011 Water Quality Results - North Basin

Sample Year Mean	Secchi Disc Depth (Metres)	Total Phosphorus Euphotic Zone (Micrograms/litre)	Total Phosphorus 1 Metre off Bottom (Micrograms/litre)	Chlorophyll <u>a</u> Composite (Micrograms/litre)
2001	3.4	12.9	9	4.91
2006	4.2	10	20	3.07
2011	4	20	14.3	4.6
N Mean	3 3.9	3 14.3	3 14.4	3 4.19
Standard Deviation	0.4163332	5.144900388	5.501211988	0.985105747

*Mean based on less than 6 measurements **Includes Recreational Lakes Program Data.

Chlorophyll a data prior to 1985 has been adjusted to reflect new lab procedures
 in filtering resulting in an increase in chl.a concentrations by 35%




Secchi Disc Measurements

A *Secchi Disc* is a black and white coloured disc used to determine water clarity. The disc is lowered into the water. The point, at which you can no longer distinguish the black and white, is called the Secchi depth. The higher the Secchi Disc measurement, the more clear your lake.

Lakes are classified as *oligotrophic*, *mesotrophic*, or *eutrophic* depending on age and whether they have little, some, or a lot of life, respectively. Oligotrophic lakes are the youngest and, usually, least fertile lakes; they tend to be deep with sparse aquatic vegetation and few fish. Mesotrophic lakes are middle-aged lakes that are less deep and more fertile than oligotrophic lakes. And eutrophic lakes (the oldest lakes) are most fertile and even more shallow than mesotrophic lakes.

Eutrophic lakes eventually reach the point where demand for oxygen exceeds the oxygen supply. Eutrophic lakes have many aquatic life forms that eventually die and decompose; [decomposition](#) uses up oxygen that could have supported additional life. Decomposing material, detritus, collects on the lake's bottom making the lake shallower. As oxygen becomes sparse, lakes approach *senescence*, full maturity to death.

Interpreting SECCHI DISC Results	
Secchi Reading	Lake Nutrient Status
Over 5 metres	Oligotrophic - unenriched, few nutrients
3.0 to 4.9 metres	Mesotrophic – moderately enriched, some nutrients
Less than 2.9 metres	Eutrophic – enriched, higher levels of nutrients



Interpreting Total Phosphorus Results


Phosphorus is the nutrient that controls the growth of algae in most Ontario lakes. For this reason any increase in phosphorus in the lake will increase the quantity of algae that can grow. High levels of phosphorus can lead to algal blooms and in some cases affect the habitat of cold water fish such as lake trout. A general guideline exists to characterize your lake based on the total phosphorus that is measured.




The Kremmerer Bottle (pictured above) is used for sampling water at specific depths. The bottle is lowered with both ends open to the required depth, a weight (on the rope) is dropped, when the weight hits the bottle the it causes both ends to close sealing the sample water in the bottle.


Interpreting TOTAL PHOSPHORUS Results	
Total Phosphorus	Lake Nutrient Status
10 ug/L or less	Oligotrophic - unenriched, few nutrients
11 to 20 ug/L	Mesotrophic – moderately enriched, some nutrients
21 ug/L or more	Eutrophic – enriched, higher levels of nutrients

Bennett Lake—South Basin Annual Mean Total Phosphorus Results (ug/L)



Interpreting TOTAL PHOSPHORUS Results (cont'd)	
Total Phosphorus	Lake Nutrient Status
10 ug/L or less	Oligotrophic - unenriched, few nutrients
11 to 20 ug/L	Mesotrophic – moderately enriched, some nutrients
21 ug/L or more	Eutrophic – enriched, higher levels of nutrients






Evaluating Chlorophyll a Results:

Water clarity is influenced by the amount of phytoplankton or microscopic algae present in the water. Chlorophyll a is the green pigment in phytoplankton. The lower the chlorophyll a density in your lake, the clearer your lake is. Chlorophyll a is directly affected by the amount of total phosphorus in your lake. The more phosphorus there is in the water, the more algal growth will occur.

The Composite Sampler (pictured above) is used for water samples similar to a core sample. The tin is dropped into the water. When it reached the required depth it is slowly pulled back to the surface. The tube is filled as water enters one tube while air escapes the other. It is essential that some air is left in the tin to ensure collection throughout the entire haul to the surface.

Interpreting CHLOROPHYLL <u>A</u> Results	
Chlorophyll <u>a</u> Reading	Lake Nutrient Status
Up to 2 ug/L - low algal density	Oligotrophic - unenriched, few nutrients
2-4 ug/L - moderate algal density	Mesotrophic - moderately enriched, some nutrients
More than 4 ug/L- high algal density	Eutrophic - enriched, higher levels of nutrients

**Bennett Lake—South and North Basins
Annual Mean Chlorophyll a Results (ug/L)**





Help MVC and the Ontario Federation of Anglers and Hunters (OFAH) Stop the Invasion!

Bennett Lake was also tested for invasive species in 2011, in particular, for zebra mussels and spiny water flea, in partnership with the Ontario Federation of Anglers and Hunters (OFAH). Bennett Lake did *not* have spiny water flea present however; zebra mussel veligers (larvae) were present in the sample collected.


Residents and property owners need to ensure that all access points to the lake have posted signs indicating the precautions they can take to avoid the spread of invasive species into Bennett Lake. Another recommendation is for residents to begin participation in the invasive species monitoring program through MVC.

The plankton haul net (pictured above) looks like a windsock with a plastic cup attached to the end. The mesh size of the net is 63 microns, which can filter microscopic organisms (plankton) like the spiny water flea and the zebra mussel veligers from the water. The plastic cup portion of the net is called the cod end and it collects the plankton sample as the water passes through the net.

Check and clean watercraft every time it is moved to a different water body

Evaluating pH Results

Lakes with pH levels at 7.3 or higher are vulnerable to zebra mussels invasive.



Call MVC at (613) 259-2421 or the Invading Species Hotline 1-800-563-7711

Sampling Results—North Basin

Bennett Lake—North Basin

#11-08

May 25, 2011

10:15 AM

Depth (Metres)	Temperature (Degrees Celsius)	Dissolved Oxygen (Milligrams / Litre)	Percent % Saturation	Thermal Stratification
0.1	17.4	9.25	96.4	Epilimnion
1	16.8	9.25	95.3	
2	16.8	9.03	93.1	
3	16.7	8.93	91.8	
4	16.6	8.83	90.8	
5	16.2	8.62	87.8	
6	14.9	2.65	75.5	Thermocline
7	Bottom	Bottom	Bottom	

Bennett Lake—North Basin

#11-08

July 27, 2011

10:15 AM

Depth (Metres)	Temperature (Degrees Celsius)	Dissolved Oxygen (Milligrams / Litre)	Percent % Saturation	Thermal Stratification
0.1	25.36	7.34	89.1	Epilimnion
1	25.33	7.17	87.3	
2	25.32	7.16	87.1	
3	25.31	7.15	87.2	
4	25.30	7.14	87.0	
5	25.28	7.08	86.0	
6	22.09	1.89	16.7	Thermocline
7	19.17	0.43	4.4	
8	Bottom	Bottom	Bottom	

Bennett Lake—North Basin

#11-08

August 31, 2011

10:25 AM

Depth (Metres)	Temperature (Degrees Celsius)	Dissolved Oxygen (Milligrams / Litre)	Percent % Saturation	Thermal Stratification
0.1	17.4	9.25	96.4	Epilimnion
1	16.8	9.25	95.3	
2	16.8	9.03	93.1	
3	16.7	8.93	91.8	
4	16.6	8.83	90.8	
5	16.2	8.62	87.8	
6	14.9	2.65	75.5	Thermocline
7	Bottom	Bottom	Bottom	

 Warm Water Fisheries Habitat (Bass, Walleye, Pike, Perch) = DO > 4 mg/L at < 25°C

Sampling Results—South Basin

Bennett Lake—South Basin

11-09

May 25, 2011

11:15 AM

Depth (Metres)	Temperature (Degree Celsius)	Dissolved Oxygen (Milligrams / Litre)	Percent % Saturation	Thermal Stratification
0.1	18.8	9.22	97.7	Epilimnion
1	17.8	9.22	96.9	
2	17.6	9.21	96.5	
3	17.5	9.23	96.5	
4	16.4	8.95	90.5	
5	14.3	8.57	83.6	Thermocline 1
6	13.5	8.38	80.4	Thermocline 2
7	12.7	8.20	77.1	
8	11.9	7.25	66.9	
9	10.4	4.26	38.0	Thermocline 3
10	9.7	3.2	28.1	Hypolimnion
11	9.2	2.67	23.2	
12	Bottom	Bottom	Bottom	

Bennett Lake—South Basin

11-09

July 27, 2011

10:59 AM

Depth (Metres)	Temperature (Degree Celsius)	Dissolved Oxygen (Milligrams / Litre)	Percent % Saturation	Thermal Stratification
0.1	25.19	8.28	100.8	Epilimnion
1	25.28	8.18	99.7	
2	25.23	8.08	98.4	
3	25.19	7.97	97.2	
4	24.18	7.32	88.1	
5	18.16	1.89	20.4	Thermocline
6	14.65	0.68	6.8	
7	12.2	0.54	5.	
8	11.12	0.48	4.5	
9	10.51	0.41	3.7	Hypolimnion
10	10.08	0.36	3.2	
11	Bottom	Bottom	Bottom	

 Warm Water Fisheries Habitat (Bass, Walleye, Pike, Perch) = DO > 4 mg/L at < 25°C

Sampling Results—South Basin (con't)

Bennett Lake—South Basin

11-09

August 31, 2011

11:42 AM

Depth (Metres)	Temperature (Degree Celsius)	Dissolved Oxygen (Milligrams / Litre)	Percent % Saturation	Thermal Stratification
0.1	22.0	8.53	97.5	Epilimnion
1	21.5	8.61	97.5	
2	21.4	8.52	96.2	
3	21.3	8.44	95.2	
4	21.2	8.21	92.4	
5	21.0	8.16	91.5	
6	20.9	7.62	85.2	
7	18.5	2.60	27.5	Thermocline
8	13.3	0.26	2.4	
9	11.8	0.08	0.7	
10	10.6	0.02	0.2	
11	10.4	-0.02	-0.2	Hypolimnion
12	10.3	-0.04	-0.3	
13	Bottom	Bottom	Bottom	



Warm Water Fisheries Habitat (Bass, Walleye, Pike, Perch) = DO > 4 mg/L at < 25°C

Seine Netting

Seine netting by hand is a way of sampling fish species that may live or visit the near shore areas of a waterbody. A seine net is a type of fishing net that has floats along one edge and weight along the other edge, to keep it upright in the water. It is then dragged through a section of water, encircling it, thus collecting all the fish within that area. The depth of the testing area is limited to areas wadeable by the field crew. This method has a very limited impact on the health of the fish sampled and is affordable, easy to do, and portable.



Seine netting was conducted at the boat launch sites of all the 2011 watershed watch lakes to help expand our knowledge of each lake beyond just its chemistry. Netting was conducted in July and August to avoid disturbing sensitive nesting, and breeding sites. A good variety of game fish and bait fish were found this year and the species found at Bennett Lake can be found in the chart below.

The majority of the individual fish captured with the seine net are bait fish such as minnows and cyprinids; some juvenile and adult game fish were also caught. Both groups (bait fish and juvenile game fish) tend to stick close to shore to avoid predation from larger fish that can be found in deeper waters. Near shore areas may also contain aquatic vegetation which is ideal camouflage for all sizes of fish that are either hiding from predators, or waiting to surprise prey, explaining why some adult game fish were caught.

It is important to note that if something was not caught in the seine we cannot conclude that the fish species is not in the lake rather that the species was not in the sampling zone when the sampling was done.

Most stakeholder interest in fish species within a water body has to do with game fish. However, baitfish far outnumber game fish and thus play a critical role within their ecosystem and the food chain. It is important to take note of their presence, and provide them the same consideration you would for larger fish. If you are curious about learning more about baitfish, and how to identify the different species, please refer to the Department of Fisheries and Oceans "[Baitfish Primer](#)", it is available online and at the MVC office.

Lake	Fish Species Caught
Bennett Lake	Pumpkinseed
	Largemouth Bass
	Banded Killifish
	Spottail shiner
	Yellow Perch
	Black Crappie



MVC would like to thank the many dedicated volunteers and the Lake Steward Network for their assistance with and support of the Watershed Watch program.

For more information about the MVC Monitoring Program please call: 613.259.2421 ext. 235 or email: monitoring@mvc.on.ca or visit: www.mvc.on.ca

We are located at
4175 Hwy 511
Lanark Ontario
K0G 1K0



Mississippi Valley Conservation

community • environment • balance